



TITLE:

# COMMENTS OF MULTILINEAR STRONG MAXIMAL OPERATORS ON MIXED LEBESGUE SPACES (The deepening of function spaces and its environment)

AUTHOR(S):

Yabuta, Kôzô

---

CITATION:

Yabuta, Kôzô. COMMENTS OF MULTILINEAR STRONG MAXIMAL OPERATORS ON MIXED LEBESGUE SPACES (The deepening of function spaces and its environment). 数理解析研究所講究録 2018, 2095: 57-63

ISSUE DATE:

2018-12

URL:

<http://hdl.handle.net/2433/251705>

RIGHT:

# COMMENTS ON MULTILINEAR STRONG MAXIMAL OPERATORS ON MIXED LEBESGUE SPACES

KÔZÔ YABUTA

**0.1. Maximal type operators.** Let  $n \geq 1$  and  $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ . Let  $\mathcal{M}$  be the well-known Hardy-Littlewood maximal operator defined on  $\mathbb{R}^n$  as follows.

$$\mathcal{M}f(x) = \sup_{r>0} \frac{1}{|B_r(x)|} \int_{B_r(x)} |f(y)| dy,$$

where  $B_r(x)$  is the open ball in  $\mathbb{R}^n$  centered at  $x$  with radius  $r$  and  $|B_r(x)|$  denotes the volume of  $B_r(x)$ . The corresponding uncentered maximal function will be denoted by  $\widetilde{\mathcal{M}}f$ .

It is well known that  $\mathcal{M}$  and  $\widetilde{\mathcal{M}}$  are of type  $(p, p)$  for  $1 < p \leq \infty$  and weak type  $(1, 1)$ . In 1997, Kinnunen [18] first showed that  $\mathcal{M}$  is bounded from the first order Sobolev spaces  $W^{1,p}(\mathbb{R}^n)$  to  $W^{1,p}(\mathbb{R}^n)$  for  $1 < p \leq \infty$ . Later on, the  $W^{1,p}$ -bounds for the uncentered operator  $\widetilde{\mathcal{M}}$  was obtained by Hajlasz and Onninen [15]. Since then, many works have been done to extend the above results in a more general setting. We refer the reader to see [19], [20], [7, 28], [27], [31], [32]. However, the results for  $p = 1$  are quite different. In 2004, Hajlasz and Onninen [15] surprisingly pointed out that the Hardy-Littlewood maximal operator is not bounded on  $W^{1,1}$  space. Thus, it is quite natural to consider whether the operator  $f \mapsto |\nabla \mathcal{M}f|$  is bounded from  $W^{1,1}(\mathbb{R}^n)$  to  $L^1(\mathbb{R}^n)$  or not. When  $n = 1$ , this problem was completely solved in [2, 24, 26, 39]. But for  $n \geq 2$ , only partial results were given by Hajlasz and Malý [14] and Luiro [33]. For more previous works or related topic we refer the readers to consult [3, 5, 6, 8, 24, 21, 22, 29], and the references therein.

In order to study the multilinear singular integral operators with multiple weights, in 2009, Lerner et al [25] introduce the multilinear version of Hardy-Littlewood maximal functions. In 2011, Grafakos, Liu, Pérez and Torres [13] introduced and studied the weighted strong and endpoint estimates for the following multilinear strong maximal function  $\mathcal{M}_{\mathcal{R}}$ .

**Definition 0.1** (Multilinear strong maximal function [13]). Let  $\vec{f} = (f_1, \dots, f_m)$  be an  $m$ -dimensional vector of locally integrable functions. Define the multilinear strong maximal function  $\mathcal{M}_{\mathcal{R}}$  by

$$(0.1) \quad \mathcal{M}_{\mathcal{R}}(\vec{f})(x) = \sup_{\substack{R \ni x \\ R \in \mathcal{R}}} \prod_{i=1}^m \frac{1}{|R|} \int_R |f_i(y_i)| dy_i,$$

where  $x \in \mathbb{R}^n$  and  $\mathcal{R}$  denotes the family of all rectangles in  $\mathbb{R}^n$  with sides parallel to the axes.

---

*Date:* 2018.2.8, from a joint work with Feng Liu (Shandong University of Science and Technology, China), Rodolfo H. Torres (The University of Kansas, USA) and Qingying Xue (Beijing Normal University, China ).

Whenever  $m = 1$ , we simply denote  $\mathcal{M}_{\mathcal{R}}$  by  $\mathcal{M}_{\mathcal{R}}$ . Then  $\mathcal{M}_{\mathcal{R}}$  coincides with the classical strong maximal operator studied by Jessen, Marcinkiewicz and Zygmund [17] in 1935. Unlike the Hardy-Littlewood maximal function, the strong maximal function  $\mathcal{M}_{\mathcal{R}}$  is not of weak type  $(1, 1)$ . Therefore, as a replacement, Jessen, Marcinkiewicz and Zygmund [17] showed that it is bounded from  $L(\log^+ L)(\mathbb{R}^d)$  to  $L^1(\mathbb{R}^d)$ . In 1975, the  $L(\log^+ L)(\mathbb{R}^d)$  type estimate was again proved by Córdoba and Fefferman [9] by using an alternative geometric method [9].

It is known that [13]  $\mathcal{M}_{\mathcal{R}}$  is bounded from  $L^{p_1}(\mathbb{R}^d) \times \cdots \times L^{p_m}(\mathbb{R}^d)$  to  $L^p(\mathbb{R}^d)$  for all  $1 < p_1, \dots, p_m, p \leq \infty$  and  $1/p = \sum_{i=1}^m 1/p_i$ . Moreover, for each  $f_i \in L^{p_i}(\mathbb{R}^d)$ , it holds that

$$(0.2) \quad \|\mathcal{M}_{\mathcal{R}}(\vec{f})\|_{L^p(\mathbb{R}^d)} \lesssim_{p_1, \dots, p_m} \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R}^d)}.$$

**0.2. Mixed Lebesgue spaces.** We first introduce the definition of mixed Lebesgue spaces.

**Definition 0.2** (Mixed Lebesgue space  $L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$ , [23]). Let  $1 \leq p, q < \infty$  and  $n_i \geq 1$  ( $i = 1, 2$ ), the mixed Lebesgue space  $L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$  is defined by

$$L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}) := \{f : \mathbb{R}^{n_1+n_2} \rightarrow \mathbb{R}; \quad \|f\|_{L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})} < \infty\},$$

where

$$\|f\|_{L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})} := \left( \int_{\mathbb{R}^{n_1}} \left( \int_{\mathbb{R}^{n_2}} |f(x, y)|^q dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}}.$$

Similarly, we can define mixed Lebesgue space with  $\ell$  terms by  $L^{p_1, p_2, \dots, p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \cdots \times \mathbb{R}^{n_\ell})$ . It is easy to see that  $L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}) = L^p(\mathbb{R}^{n_1+n_2})$  if  $p = q$ .

It is well known that this space still preserves the following properties: it is a Banach space and some classical theorems, such like monotone convergence theorem, Lebesgue's dominated convergence theorem still hold. The definition of mixed Lebesgue space can be traced back to the nice work of Benedek and Panzone [4], in 1961. Since then, achievements have been made in the study of some classical operators on mixed Lebesgue spaces. Among these achievements, there are nice works of Adams and Bagby [1], Lizorkin [34], and Milman [35]. These works mainly focused on the translation-dilation invariant estimates for Riesz potentials, the multipliers of Fourier integrals and bounds of convolutions, and the interpolation problem of Banach spaces and Lorentz spaces with mixed norms. Later on, the maximal inequalities and Fourier multipliers for spaces with mixed quasi-norms were studied by Schmeisser [37] and the theory of vector-valued singular operators on mixed Lebesgue spaces was considered by Fernandez [11]. Recently, some weighted theory for maximal operators associated with some special rectangles constructed by the products of two cubes was developed by Kurtz [23]. Still more recently, Radial multipliers and restriction to surfaces of the Fourier transform in mixed-norm spaces were demonstrated by Córdoba and Latorre Crespo [10]. Moreover, the smoothing properties of bilinear operators and Leibniz-type rules in mixed Lebesgue spaces were presented very recently by Hart, Torres and Wu [16].

Based on the previous results for the Hardy-Littlewood maximal operators, the multilinear strong maximal functions and some other classical operators, it is therefore a quite natural question to ask whether the multilinear strong maximal operators are still

bounded on the product of mixed Lebesgue spaces. and enjoy the regularity and continuity properties.

0.3. We will consider the boundedness, regularity and continuity properties of the multilinear strong maximal operators on the mixed Lebesgue spaces and mixed Lebesgue-Sobolev spaces. We will see that these results rely heavily on one dimensional results. To begin with, we introduce the definition of mixed Lebesgue-Sobolev spaces.

**Definition 0.3 (Mixed Lebesgue-Sobolev spaces  $W^{1,p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$ ).** Let  $1 \leq p \leq \infty$ ,  $1 \leq q \leq \infty$  and  $n_i \geq 1$  ( $i = 1, 2$ ), the mixed Sobolev Lebesgue space  $W^{1,p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})$  is defined by

$$W^{1,p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}) := \{f : \mathbb{R}^{n_1+n_2} \rightarrow \mathbb{R}; \quad \|f\|_{W^{1,p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})} < \infty\},$$

where  $\|f\|_{W^{1,p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})} := \|\nabla f\|_{L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})} + \|f\|_{L^{p,q}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})}$ . Similarly, we can define mixed Lebesgue-Sobolev spaces with  $\ell$  terms by  $W^{1,p_1,p_2,\dots,p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$ .

We obtain the following boundedness of  $\mathcal{M}_{\mathcal{R}}$  on mixed Lebesgue spaces.

**Theorem 0.1 (Estimates on mixed Lebesgue spaces).** Let  $\ell \geq 1$  and  $n = \sum_{i=1}^{\ell} n_i$  with  $n_i \geq 1$  ( $i = 1, \dots, \ell$ ). Let  $1 < p_j, p_{1j}, p_{2j}, \dots, p_{mj} < \infty$  and  $1/p_j = \sum_{i=1}^m 1/p_{ij}$  ( $j = 1, \dots, \ell$ ). Then, the following inequality holds

$$\|\mathcal{M}_{\mathcal{R}}(\vec{f})\|_{L^{p_1,p_2,\dots,p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})} \lesssim_{n,p_1,\dots,p_\ell} \prod_{j=1}^m \|f_j\|_{L^{p_{j1},p_{j2},\dots,p_{j\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})}.$$

Now, we may consider the properties of  $\mathcal{M}_{\mathcal{R}}$  on mixed Lebesgue-Sobolev spaces.

**Theorem 0.2 (Estimates on mixed Lebesgue-Sobolev spaces).** Let  $\ell \geq 1$  and  $n = \sum_{i=1}^{\ell} n_i$  with  $n_i \geq 1$  ( $i = 1, \dots, \ell$ ). Let  $1 < p_j, p_{1j}, p_{2j}, \dots, p_{mj} < \infty$  and  $1/p_j = \sum_{i=1}^m 1/p_{ij}$  ( $j = 1, \dots, \ell$ ). Then  $\mathcal{M}_{\mathcal{R}}$  is bounded from products of Lebesgue-Sobolev spaces  $W^{1,p_{11},p_{12},\dots,p_{1\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell}) \times \dots \times W^{1,p_{m1},p_{m2},\dots,p_{m\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$  to Lebesgue-Sobolev spaces  $W^{1,p_1,p_2,\dots,p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$ , and it holds that

$$\|\mathcal{M}_{\mathcal{R}}(\vec{f})\|_{W^{1,p_1,p_2,\dots,p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})} \lesssim_{n,p_1,\dots,p_\ell} \prod_{j=1}^m \|f_j\|_{W^{1,p_{j1},p_{j2},\dots,p_{j\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})}$$

Moreover, the following property holds

$$|D_l \mathcal{M}_{\mathcal{R}}(\vec{f})(x)| \leq \sum_{j=1}^m \mathcal{M}_{\mathcal{R}}(\vec{F}_j)(x), \quad \text{a.e. } x \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell},$$

where  $\vec{F}_j = (f_1, \dots, f_{j-1}, D_l f_j, f_{j+1}, \dots, f_m)$ .

**Theorem 0.3 (Continuity on mixed Lebesgue-Sobolev spaces).** Let  $\ell \geq 1$  and  $n = \sum_{i=1}^{\ell} n_i$  with  $n_i \geq 1$  ( $i = 1, \dots, \ell$ ). Let  $1 < p_j, p_{1j}, p_{2j}, \dots, p_{mj} < \infty$ ,  $p_{1j} \geq p_{2j} \geq \dots \geq p_{mj} > 1$  and  $1/p_j = \sum_{i=1}^m 1/p_{ij}$  ( $j = 1, \dots, \ell$ ). Then  $\mathcal{M}_{\mathcal{R}}$  is continuous from products of Lebesgue-Sobolev spaces  $W^{1,p_{11},p_{12},\dots,p_{1\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell}) \times \dots \times W^{1,p_{m1},p_{m2},\dots,p_{m\ell}}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$  to Lebesgue-Sobolev spaces  $W^{1,p_1,p_2,\dots,p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$ .

**Remark 0.4.** If  $f_i, \nabla f_i \in L^r(\mathbb{R}^n)$  ( $i = 1, 2, \dots, m$ ) for some  $1 < r < \infty$ , then the assumption  $p_{1j} \geq p_{2j} \geq \dots \geq p_{mj} > 1$  can be removed in Theorem 0.3.

We state here some comments on fundamental facts on mixed  $L^p$  spaces, used in the proofs of the above results.

**Lemma 0.4.** *Let  $1 \leq p_2 \leq p_1 < \infty$  and  $n_1, n_2 \in \mathbb{N}$ . Then*

$$\|f\|_{L^{p_2, p_1}(\mathbb{R}^{n_2} \times \mathbb{R}^{n_1})} \leq \|f\|_{L^{p_1, p_2}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})}.$$

*Proof.* By Minkowski's inequality

$$\begin{aligned} \|f\|_{L^{p_2, p_1}(\mathbb{R}^{n_2} \times \mathbb{R}^{n_1})} &= \left( \int_{\mathbb{R}^{n_1}} \left( \int_{\mathbb{R}^{n_2}} |f(x_1, x_2)|^{p_2} dx_2 \right)^{p_1/p_2} dx_1 \right)^{1/p_1} \\ &\leq \left( \int_{\mathbb{R}^{n_2}} \left( \int_{\mathbb{R}^{n_1}} |f(x_1, x_2)|^{p_1} dx_1 \right)^{p_2/p_1} dx_2 \right)^{1/p_2} \\ &= \|f\|_{L^{p_1, p_2}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2})}. \end{aligned}$$

□

From this, it follows immediately the following:

**Lemma 0.5.** *Let  $1 \leq p_1, p_2, \dots, p_\ell < \infty$ ,  $p_j \leq p_{j-1}$  and  $n_1, n_2, \dots, n_\ell \in \mathbb{N}$ . Then,*

$$\|f\|_{L^{p_1, \dots, p_{j-2}, p_j, p_{j-1}, p_{j+1}, \dots, p_\ell}(\mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_{j-2}} \times \mathbb{R}^{n_j} \times \mathbb{R}^{n_{j-1}} \times \mathbb{R}^{n_{j+2}} \times \dots \times \mathbb{R}^{n_\ell})} \leq \|f\|_{L^{p_1, p_2, \dots, p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})}.$$

Using this we get

**Lemma 0.6.** *Let  $1 \leq p_\ell \leq p_{\ell-1} \leq \dots \leq p_1 < \infty$ , and  $n_1, n_2, \dots, n_\ell \in \mathbb{N}$ . Then, for any  $2 \leq j \leq \ell$*

$$\|f\|_{L^{p_j, p_1, \dots, p_{j-1}, p_{j+1}, \dots, p_\ell}(\mathbb{R}^{n_j} \times \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_{j-1}} \times \mathbb{R}^{n_{j+2}} \times \dots \times \mathbb{R}^{n_\ell})} \leq \|f\|_{L^{p_1, p_2, \dots, p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})}.$$

From this it follows

**Remark 0.1.** Under the assumption of Lemma 0.6, for  $f \in L^{p_1, p_2, \dots, p_\ell}(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \dots \times \mathbb{R}^{n_\ell})$  we have

$$f(x_1, \dots, x_{j-1}, \cdot, x_{-j+1}, \dots, x_\ell) \in L^{p_j}(\mathbb{R}^{n_j})$$

for almost every  $(x_1, \dots, x_{j-1}, x_{-j+1}, \dots, x_\ell) \in \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_{j-1}} \times \mathbb{R}^{n_{j+2}} \times \dots \times \mathbb{R}^{n_\ell}$ .

**Remark 0.2.** Without the assumption of Lemma 0.6, the conclusion in the above remark does not hold. In fact, let  $1 \leq p < q < \infty$ . Taking  $1/q < a < 1/p$ , set

$$f(x) = \chi_{-|x| < 1, |y| < 1} \frac{1}{|x - y|^a}.$$

Then  $f \in L^{p, q}(\mathbb{R} \times \mathbb{R})$  but  $f(x, y) \notin L^q(\mathbb{R}_y)$  for  $|x| < 1$ , and hence  $f \notin L^{q, p}(\mathbb{R} \times \mathbb{R})$ .

*Proof.*

$$\begin{aligned} \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)|^p dx \right)^{q/p} dy \right)^{1/q} &= \left( \int_{|y| < 1} \left( \int_{|x| < 1} \frac{1}{|x - y|^{ap}} dx \right)^{q/p} dy \right)^{1/q} \\ &\leq \left( \int_{|y| < 1} \left( \int_0^1 \frac{2}{x^{ap}} dx \right)^{q/p} dy \right)^{1/q} \\ &= 2^{1/q} \left( \int_0^1 \frac{2}{x^{ap}} dx \right)^{1/p} < \infty. \end{aligned}$$

But

$$\int_{\mathbb{R}} |f(x, y)|^q dy = \int_{|y| < 1} \frac{dy}{|x - y|^{aq}} = +\infty \quad \text{for } |x| < 1,$$

and hence  $f \notin L^{q, p}(\mathbb{R} \times \mathbb{R})$ . □

**Example 0.5.** Let  $1 \leq p < q < \infty$  and  $1/q < \alpha < 1/p$ . Set

$$f(x, y) = \chi_{\{|x-y| \leq 1\}} \frac{1}{(|x-y|(1+|y|))^\alpha}.$$

Then  $f \in L^{p,q}(\mathbb{R} \times \mathbb{R})$ , but  $f(x, \cdot) \notin L^q(\mathbb{R}_y)$  for any  $x \in \mathbb{R}$ . In particular,  $f \notin L^{q,p}(\mathbb{R} \times \mathbb{R})$ .

Furthermore,  $f \notin L^r(\mathbb{R}^2)$  for any  $1 \leq r < \infty$ .

*Proof.*

$$\begin{aligned} \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)|^p dx \right)^{q/p} dy \right)^{1/q} &= \left( \int_{\mathbb{R}} \left( \int_{|x-y| \leq 1} \frac{1}{|x-y|^{p\alpha}} dx \right)^{q/p} \frac{1}{(1+|y|)^{q\alpha}} dy \right)^{1/q} \\ &\leq \left( \int_{\mathbb{R}} \left( \int_0^1 \frac{2}{x^{p\alpha}} dx \right)^{q/p} \frac{1}{(1+|y|)^{q\alpha}} dy \right)^{1/q} \\ &= \left( \frac{2}{1-p\alpha} \right)^{1/p} \left( \frac{2}{q\alpha-1} \right)^{1/q} < \infty. \end{aligned}$$

On the other hand, for  $|x-y| \leq 1$ , we have  $|y| \leq |x| + |y-x| \leq |x| + 1$ . So, we get

$$\left( \int_{\mathbb{R}} |f(x, y)|^q dy \right)^{1/q} = \left( \int_{|x-y| \leq 1} \frac{1}{(|x-y|(1+|y|))^{q\alpha}} dy \right)^{1/q} \geq \left( \int_{|y| \leq 1} \frac{dy}{|y|^{q\alpha}} \right)^{1/q} \cdot \frac{1}{(|x|+2)^\alpha} = +\infty.$$

which implies  $f(x, \cdot) \notin L^q(\mathbb{R}_y)$  for any  $x \in \mathbb{R}$ .

Next, for  $1/\alpha \leq r < \infty$  we have as above

$$\left( \int_{\mathbb{R}} |f(x, y)|^r dy \right)^{1/r} = \left( \int_{|x-y| \leq 1} \frac{1}{(|x-y|(1+|y|))^{r\alpha}} dy \right)^{1/r} \geq \left( \int_{|y| \leq 1} \frac{dy}{|y|^{r\alpha}} \right)^{1/r} \cdot \frac{1}{(|x|+2)^\alpha} = +\infty.$$

which implies  $f(x, \cdot) \notin L^r(\mathbb{R}_y)$  for any  $x \in \mathbb{R}$ , and so  $f(x, y) \notin L^r(\mathbb{R}^2)$ .

If  $1 \leq r < 1/\alpha$ , we get

$$\begin{aligned} \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)|^r dx \right) dy \right)^{1/r} &= \left( \int_{\mathbb{R}} \left( \int_{|x-y| \leq 1} \frac{1}{|x-y|^{r\alpha}} dx \right) \frac{1}{(1+|y|)^{r\alpha}} dy \right)^{1/r} \\ &= \left( \int_{\mathbb{R}} \left( \int_0^1 \frac{2}{x^{r\alpha}} dx \right) \frac{1}{(1+|y|)^{r\alpha}} dy \right)^{1/r} \\ &= \left( \frac{2}{1-r\alpha} \right)^{1/r} \left( \int_{\mathbb{R}} \frac{1}{(1+|y|)^{r\alpha}} dy \right)^{1/r} = \infty, \end{aligned}$$

i.e.  $f \notin L^r(\mathbb{R}^2)$ . □

## REFERENCES

- [1] D. R. Adams and R. J. Bagby, *Translation-dilation invariant estimates for Riesz potentials*, Indiana Univ. Mat. J. **23** (1974), 1051-1067.
- [2] J. M. Aldaz and J. Pérez Lázaro, *Functions of bounded variation, the derivative of the one dimensional maximal function, and applications to inequalities*, Trans. Amer. Math. Soc., **359** (5) (2007), 2443-2461.
- [3] J. M. Aldaz and J. Pérez Lázaro, *Boundedness and unboundedness results for some maximal operators on functions of bounded variation*, J. Math. Anal. Appl., **337** (2008), 130-143.
- [4] A. Benedek and R. Panzone, *The spaces  $W$  with mixed norm*, Duke Math. J., **28** (1961), 301-324.
- [5] E. Carneiro and J. Mardid, *Derivative bounds for fractional maximal functions*, Trans. Amer. Math. Soc., **369** (2017), 4063-4092.

- [6] E. Carneiro, J. Mardid, L. B. Pierce, *Endpoint Sobolev and BV continuity for maximal operators*, J. Funct. Anal., **273**(10), 3262-3294(2017)
- [7] E. Carneiro and D. Moreira, *On the regularity of maximal operators*, Proc. Amer. Math. Soc., **136** (12) (2008), 4395-4404.
- [8] E. Carneiro and B. F. Svaiter, *On the variation of maximal operators of convolution type*, J. Funct. Anal., **265** (2013), 837-865.
- [9] A. Córdoba and R. Fefferman, *A geometric proof of the strong maximal theorem*, Ann. Math., (2) **102** (1) (1975), 91-95.
- [10] A. Córdoba, E. Latorre Crespo, *Radial multipliers and restriction to surfaces of the Fourier transform in mixed-norm spaces*, Math. Z., **286** no. 3-4 (2017), 1479-1493.
- [11] D. L. Fernandez, *Vector-valued singular operators on  $L^p$ -spaces with mixed norms and applications*, Pacific J. Math., **129**, No. 2 (1987), 257-275.
- [12] L. Grafakos, *Classical and Modern Fourier Analysis*, Pearson Education, Upper Saddle River, New Jersey, 2004.
- [13] L. Grafakos, L. Liu, C. Pérez and R. H. Torres, *The multilinear strong maximal function*, J. Geom. Anal., **21**, (2011), 118-149.
- [14] Hajlasz, P, Malý, J: *On approximate differentiability of the maximal function*, Proc. Amer. Math. Soc., **138**(1), 165-174(2010)
- [15] P. Hajlasz and J. Onninen, *On boundedness of maximal functions in Sobolev spaces*, Ann. Acad. Sci. Fenn. Math., **29** (1) (2004), 167-176.
- [16] J. Hart, R. H. Torres, X. Wu, *Smoothing properties of bilinear operators and Leibniz-type Rules in Lebesgue and mixed Lebesgue spaces*, <https://arxiv.org/abs/1701.02631>.
- [17] B. Jessen, J. Marcinkiewicz and A. Zygmund, *Note on the differentiability of multiple integrals*, Fund. Math., **25** (1935), 217-234.
- [18] J. Kinnunen, *The Hardy-Littlewood maximal function of a Sobolev function*, Israel J. Math., **100** (1997), 117-124.
- [19] J. Kinnunen and P. Lindqvist, *The derivative of the maximal function*, J. Reine. Angew. Math., **503** (1998), 161-167.
- [20] J. Kinnunen and E. Saksman, *Regularity of the fractional maximal function*, Bull. London Math. Soc., **35** (4) (2003), 529-535.
- [21] S. Korry, *Boundedness of Hardy-Littlewood maximal operator in the framework of Lizorkin-Triebel spaces*, Rev. Mat. Complut., **15**(2) (2002), 401-416.
- [22] S. Korry, *A class of bounded operators on Sobolev spaces*, Arch. Math., **82** (1) 2004, 40-50.
- [23] D. S. Kurtz, *Classical operators on mixed-normed spaces with product weights*, Rocky Mountain J. Math., **37** (2007), 269-283.
- [24] O. Kurka, *On the variation of the Hardy-Littlewood maximal function*, Ann. Acad. Sci. Fenn. Math., **40** (2015), 109-133.
- [25] A. K. Lerner, S. Ombrosi, C. Pérez, R. H. Torres and R. Trujillo-González, *New maximal functions and multiple weighted for the multilinear Calderón-Zygmund theory*, Adv. Math., **220** (2009), 1222-1264.
- [26] F. Liu, T. Chen and H. Wu, *A note on the endpoint regularity of the Hardy-Littlewood maximal functions*, Bull. Austral. Math. Soc., **94** (2016), 121-130.
- [27] F. Liu and S. Mao, *On the regularity of the one-sided Hardy-Littlewood maximal functions*, Czech. Math. J., **67**(1) (2017), 219-234.
- [28] F. Liu and H. Wu, *On the regularity of the multisublinear maximal functions*, Canad. Math. Bull., **58**(4) (2015), 808-817.
- [29] F. Liu and H. Wu, *On the regularity of maximal operators supported by submanifolds*, J. Math. Anal. Appl., **453** (2017), 144-158.
- [30] F. Liu, Q. Xue and K. Yabuta, *Regularity and continuity of the strong maximal operators associated with rectangles*, <https://arxiv.org/abs/1801.09828>.
- [31] H. Luiro, *Continuity of the maximal operator in Sobolev spaces*, Proc. Amer. Math. Soc., **135**(1) (2007), 243-251.
- [32] H. Luiro, *On the regularity of the Hardy-Littlewood maximal operator on subdomains of  $\mathbb{R}^n$* , Proc. Edinburgh Math. Soc., **53**(1) (2010), 211-237.

- [33] H. Luiro, *The variation of the maximal function of a radial function*. Ark. Math., (to appear)
- [34] P. I. Lizorkin, Multipliers of Fourier integrals and bounds of convolutions in spaces with mixed norms, Izv. Akad. Nauk SSSR., 4 (1970), 225-254.
- [35] M. Milman, On interpolation of  $2^n$  Banach spaces and Lorentz spaces with mixed norms, J. Funct. Anal. **41** (1981), 1-7.
- [36] Y. Sawano, *A Handbook of Harmonic Analysis*, <http://www.comp.tmu.ac.jp/yoshihiro/teaching/harmonic-analysis/harmonic-analysis-textbook.pdf>.
- [37] H. J. Schmeisser, *Maximal inequalities and Fourier multipliers for spaces with mixed quasi-norms, Applications*, Z. Anal. Anwendungen., **3** (1984), 153-166.
- [38] B. Stöckert, *Ungleichungen vom Plancherel-Polya-Nikol'skij-Typ in gewichteten  $L_p^\Omega$ -Räumen mit gemischten Normen*, Math. Nachr., **86** (1978), 19-32.
- [39] H. Tanaka, *A remark on the derivative of the one-dimensional Hardy-Littlewood maximal function*, Bull. Austral. Math. Soc., **65**(2) (2002), 253-258.

KÔZÔ YABUTA, RESEARCH CENTER FOR MATHEMATICAL SCIENCES, KWANSEI GAKUIN UNIVERSITY, GAKUEN 2-1, SANDA 669-1337, JAPAN